

Understanding landscape resilience by changes at fine scale

Ileana Pătru-Stupariu^{1,2}, Ana-Maria Calotă³, Paulina Anastasiu⁴

1. Institute of Research of University of Bucharest, ICUB; Transdisciplinary Research Centre Landscape-Territory-Information Systems, CeLTIS, Splaiul Independentei nr. 91-95, 050095 Bucharest, Romania 2. Department of Regional Geography and Environment, Faculty of Geography, University of Bucharest, Bd. N. Bălcescu, 1, 010041 Bucharest, Romania 3. Faculty of Geography, University of Bucharest, Bd. N. Bălcescu, 1, 010041 Bucharest, Romania 4. Faculty of Biology, University of Bucharest, 1-3 Portocalelor Intr., 060101 Bucharest, Romania

Table 1 Ecological indicators according Sârbu et al. (2013)

Ecological indicator	1-3	4-6	7-X
Light	shadow	partial shadow	full light
Temperature	cold	warm	hot
Soil humidity	dry	slight moisture	aquatic
Soil reaction	acid	neutral	alkaline
Amount of N mineral in the soil	low	average	high




Figure 1 Ecological indicators for species identified near the wind farm (T0) and the corresponding sampling point on the control transect (M0)




Figure 2 Structural phytocoenosis differences between the two sites

Reference: Sârbu I., Ștefan N., Oprea A., (2013). *Plante vasculare din România, Determinator ilustrat de teren*. București: Editura VictorBVictor